

Pharmacology of angiotensin II receptors in the kidney

MARC DE GASPARO and NIGEL R. LEVENS

Cardiovascular Research Department, CIBA-GEIGY Limited, Basel, Switzerland

Pharmacology of angiotensin II receptors in the kidney. Within the kidney angiotensin II (Ang II) exerts potent effects on renal function. The intrarenal actions of Ang II include modulation of renal blood flow, glomerular filtration rate, tubular epithelial transport, renin release and cellular growth. The actions of Ang II on the kidney are mediated by specific intrarenal receptors which, based upon physical characteristics and the selective binding of non-peptide and peptide analogs may be divided into two main subtypes, termed AT₁ and AT₂. AT₁ receptors are present within the kidneys of all species and are located predominantly in the glomerulus, the renal tubules and the renal vasculature, including the afferent and efferent arterioles. Modulation of AT₁ receptors within the kidney has been shown to mediate essentially all of the known intrarenal effects of Ang II. AT₁ receptors and particularly AT₂ receptors are expressed in large numbers in fetal kidney where they may play a role in development and maturation. In some species, intrarenal AT₂ receptors disappear shortly after birth. In those species where AT₂ receptors are present in the adult kidney their role in the control of renal function has not yet been clearly defined.

Within the kidney angiotensin II (Ang II) exerts potent effects on renal function. The intrarenal actions of Ang II include modulation of renal blood flow, glomerular filtration rate, tubular epithelial transport, renin release and cellular growth. Ang II binding sites have been localized in the kidney using *in vitro* autoradiography and in many cases have been correlated to changes in physiological function [1–3]. Variations in structural activity requirements and differences in signal transduction and regulatory mechanisms in several of the intrarenal actions of Ang II have been noted over the years [3]. These observations have led to the suggestion that the actions of Ang II within the kidney are mediated by different receptor subtypes. The recent availability of new peptide and non-peptide ligands has allowed this hypothesis to be confirmed with the unambiguous characterization of at least two distinct Ang II receptor subtypes [4]. The pharmacology of Ang II receptor subtypes within the kidney are summarized in the following brief review.

Characteristics of Ang II receptor subtypes

Recently, two receptor subtypes having clearly different binding characteristics were described in aortic smooth muscle, uterus and adrenal cortex [5, 6]. These initial observations have since been confirmed in a large number of tissues including the kidney [7–10]. Ang II receptors having a high affinity for losartan but a low affinity for CGP 42112 and PD 123319 are termed AT₁; whereas,

Ang II receptors with a high affinity for CGP 42112 and PD 123319 but a low affinity for losartan are termed AT₂ [11].

The distribution of AT₁ and AT₂ receptor subtypes is heterogeneous between tissues. Some tissues, for example the liver and vascular smooth muscle cells in culture, express only the AT₁ receptor [5, 8], whereas only the AT₂ receptor is expressed in cerebellum, ovary follicles or the human uterus [5, 12, 13]. In some tissues, of which the adrenal cortex and heart are examples, both the AT₁ and AT₂ receptor subtypes are expressed [5, 7].

Three major characteristics distinguish the AT₁ and AT₂ receptor subtypes. Firstly, sulfhydryl reducing agents such as dithiothreitol decrease the binding affinity of Ang II for the AT₁ receptor, whereas they increase the affinity of Ang II for the AT₂ receptor [11]. Secondly, in contrast to the AT₁ receptor, the AT₂ receptor does not appear to be coupled to a classical G-protein [14]. Finally, the AT₁ receptor-Ang II complex is rapidly internalized within cells, whereas the complex of Ang II with the AT₂ receptor is not [15, 16].

Both the AT₁ and AT₂ receptor have been cloned and are members of the 7-transmembrane domain superfamily, but have only a 30 to 35% amino acid sequence identity [17–20]. It has been clearly established that the AT₁ receptor is coupled to phospholipase C, adenylate cyclase and voltage-dependent calcium channels [21]. In contrast, the coupling mechanism of the AT₂ receptor has not yet been clearly defined, although evidence is accumulating to suggest linkage to a protein tyrosine phosphatase [14, 20].

Ang II receptor subtypes in the kidney

Autoradiography coupled with competitive binding studies have been used to characterize the distribution of Ang II receptor subtypes in renal tissue. Using these techniques, the distribution of AT₁ and AT₂ receptor subtypes within the kidney has been shown to be species dependent. For example, in the rat and rabbit kidney Ang II receptors are essentially of the AT₁ subtype [10, 22–24] while both AT₁ and AT₂ receptors are present within the kidney of the opossum and of primates including humans [23–26]. The distribution of renal Ang II receptors has been most extensively studied in the rat where the AT₁ receptor subtype is predominantly located in the glomerulus, the renal tubules and the renal vasculature [27–29]; these observations have been confirmed using reverse transcriptase, polymerase chain reactions and *in situ* hybridization [30–32]. In these latter studies, large signals for the AT₁ receptor were detected in the glomerulus, renal papilla, proximal convoluted tubule, proximal straight tubule, cortical collecting duct and the renal vascular system. Smaller signals for the AT₁ receptor have also been detected in medullary thick ascending limb, outer collecting duct and the




Fig. 1. Effect of PD 123319 (■) and CGP 42112B (□) in comparison to losartan (○) on parameters of renal function in the sodium-depleted anesthetized rat. The results represent the changes in each parameter of renal function produced by CGP 42112B, PD 123319 and losartan relative to baseline values. Statistics compare the changes in each parameter at each dose of drug relative to control values. The methodology and detailed results of these experiments have been previously described [54]. Reprinted with permission from the *European Journal of Pharmacology* [54].




Fig. 1. Continued.

inner medullary collecting duct. Within the glomerulus, AT₁ mRNA localizes in mesangial areas, predominantly at the vascular pole and on the terminal portion of the afferent arteriole. In contrast, intense staining in all small arteries and arterioles,

including both afferent and efferent arterioles has been observed using polyclonal antisera raised against a peptide sequence of the rat AT₁ receptor [33].

In agreement with conclusions derived from autoradiography

and receptor binding studies, AT₂ mRNA has not been detected by Northern blot analysis in the rat kidney [20]. However, in the rabbit the fibrous sheath around the kidney contains an exclusive population of AT₂ receptors [34]. In contrast to the rat and the rabbit, the AT₂ receptor subtype has been demonstrated in the renal cortex of the rhesus monkey on the intrarenal vasculature and on the juxtaglomerular apparatus [23]. In human kidney, the AT₂ receptor subtype is clearly present in large preglomerular vessels of the renal cortex and in the tubulointerstitium [26, 28].

Heterogeneity of renal Ang II receptor subtypes

Only a single gene encoding for the AT₁ receptor is expressed in human kidney [35, 36]. However, in rat kidney, as well as other tissues from this species, two clones encoding for different AT₁ receptor subtypes have been identified from genomic libraries [37]. These AT₁ receptor clones termed AT_{1A} and AT_{1B} have over 90% identity but differ in their amino acid sequence, primarily in the last intracellular loop of the receptor. Both receptor subtypes have essentially similar binding signatures and are pharmacologically indistinguishable from each other [38]. AT_{1A} and AT_{1B} more accurately describe isoforms rather than distinct subtypes of the Ang II receptor. The distribution of AT_{1A} and AT_{1B} receptor isoforms within the kidney have not been extensively studied although the AT_{1A} receptor, which is expressed predominantly in vascular smooth muscle, is most abundant in the glomerular mesangial area, the vascular component of the juxtaglomerular apparatus and the terminal portion of the afferent arteriole [31].

Receptor binding studies have also suggested the existence of two AT₁ receptor subpopulations in cultured rat mesangial cells [39]. One subpopulation, which has been termed AT_{1A}, represents 86% of the total number of binding sites and has a high affinity for losartan but a low affinity for CGP 42112 and PD123319. This receptor subtype thus resembles the AT_{1A} isoform described determined using molecular biology techniques and reported above. The remaining 14% of binding sites which have unfortunately been termed AT_{1B} have a 100-fold lower affinity for losartan and a 10'000-fold higher affinity for PD 123319 [39]. This latter subtype, despite its high affinity for PD 123319, differs from the typical AT₂ receptor subtype in having a low affinity for CGP 42112B and being coupled to a G-protein. Furthermore, since the AT_{1B} receptor subtype described by receptor binding methodology has remarkably different binding properties to the AT_{1B} receptor subtype described by molecular biology techniques they must be structurally distinct. The description of subgroups of Ang II receptors within the kidney which are clearly different but with similar labeling is confusing and requires standardization.

All AT₁ subtypes in the kidney are sensitive to GTP analogues and pertussis toxin and are linked to IP₃ generation, calcium mobilization and adenylate cyclase inhibition [10, 28, 39, 40]. Mesangial AT_{1A} receptors appear to mediate Ang II induced protein synthesis [41].

Binding studies performed with radioactive losartan in isolated rat glomeruli have demonstrated that in addition to labeling the AT₁ binding site, losartan may also recognize an additional receptor with characteristics different from known AT₁ or AT₂ receptors [29]. The significance of this observation is unclear at the present time.

Renal Ang II receptors during development

Within the kidney as well as in other tissues, both AT₁ and AT₂ receptors are developmentally regulated [9, 42]. The AT₂ receptor subtype predominates in tissues of the developing rat and human fetus, where they have a predominantly glomerular and medullary distribution [43, 44]. Although less numerous, AT₁ receptors are also present in the developing rat kidney. With increasing development AT₂ receptors disappear and the AT₁ receptor increases in number to become, at least in rat kidney, the predominant receptor subtype. Since all components of the renin-angiotensin system are present within the developing fetal/placental unit and Ang II plays a role in cell growth it has been suggested that Ang II acting through AT₂ receptors and perhaps also AT₁ receptors contribute to the growth and development of the immature kidney. This hypothesis is supported by the recent observation that treatment of fetal rats with either an angiotensin converting enzyme inhibitors or an AT₁ receptor antagonists leads to abnormalities of renal growth and to persistent defects in renal function in the adult animal [45].

Changes in renal function mediated by Ang II receptors

AT₁ receptors

Consistent with the intrarenal distribution described above, AT₁ receptors have been shown to mediate essentially all of the known effects of Ang II on kidney function [46–50]. The intrarenal actions of Ang II mediated by AT₁ receptors include vasoconstriction, alterations in glomerular hemodynamics and tubular reabsorption. AT₁ receptors present upon juxtaglomerular apparatus also mediate the negative feedback of Ang II on renin release and renal cell growth [23, 51].

AT₂ receptors

In contrast to the wealth of knowledge concerning the functional role of intrarenal Ang II receptors, little is known concerning the actions of Ang II mediated through the AT₂ receptor. There have been several reports suggesting that AT₂ ligands can affect renal function. For example, the AT₂ receptor ligands PD 123177 and PD 123319 have been shown to antagonize the renal vasoconstrictor response to exogenous Ang II and to induce an increase in urine volume and free water formation in the anesthetized dog [52]. Furthermore, in the conscious rat, PD 123177 has been shown to increase renal blood flow following experimental myocardial infarction and in the anaesthetized rat to increase glomerular filtration rate, urine volume, chloride and bicarbonate excretion [46, 53]. Since in the aforementioned studies, PD 123177 and PD 123319 were used at very high doses and in the majority of cases produced effects qualitative similar to AT₁ receptor antagonists, the possibility exists that PD 123177 and PD 123319 influence renal function by interacting with the AT₁ receptor or may exhibit non-specific properties. Both PD 123177 and PD 12319 are closely related compounds, and studies of renal function with other ligands selective for the AT₂ receptors have not yet been performed. Therefore it is unknown whether the renal actions of PD 123177 and PD 123319 are specific for this class of compounds or are a property of all ligands selective for the AT₂ receptor. Recently the role of AT₂ receptors in the control of kidney function has been reassessed using the highly selective AT₂ ligand CGP 42112 [54]. In these experiments, the renal actions of CGP 42112 in comparison to PD 123319 and

losartan were investigated in sodium-depleted anesthetized rats. The results in Figure 1 illustrate that PD 123319 at infusion rates of 1 to 1000 $\mu\text{g/kg/min}$ had no effect on renal function, while CGP 42112 affected renal function only at an infusion rate of 1000 $\mu\text{g/kg/min}$. During these experiments the plasma levels of both compounds was monitored and compared to their known affinities for AT_1 and AT_2 receptors. Assuming the plasma concentrations of CGP 42112B and PD 123319 to be in equilibrium with those at the receptor level then at an infusion rates of 1 and 1000 $\mu\text{g/kg/min}$ PD 123319 (plasma levels 70 nM and 25 μM , respectively) would be expected to interact essentially with an exclusive population of AT_2 receptors. At doses between 1 and 100 $\mu\text{g/kg/min}$, CGP 42112 (plasma levels of 14 and 700 nM, respectively) would be expected to bind to AT_2 receptors but at 1000 $\mu\text{g/kg/min}$ (plasma levels 2.2 μM) to bind to 80% of the available AT_1 population. These results therefore suggest that AT_2 receptors are not involved in the control of renal function, at least in the anesthetized rat. It remains to be determined whether this is also the case in other species.

Summary

The actions of Ang II are mediated by at least two specific subtypes of receptor termed AT_1 and AT_2 . The presence or ratio of AT_1 to AT_2 receptors within the kidney varies between species. In man only one form of the AT_1 receptor subtype is expressed whereas in the rat at least three closely related isoforms are expressed. The AT_1 receptor mediates essentially all of the known effects of Ang II on the kidney. Both the AT_1 receptor and particularly the AT_2 receptor are expressed in large numbers in the fetal kidney where they may play a role in growth and differentiation. In some species AT_2 receptors disappear from the kidney shortly after birth. In those species where they remain their role in the control of renal function has not been clearly defined. Additional work is needed to define the properties, intracellular mediators and pharmacological characteristics of intrarenal Ang II receptors.

Reprint requests to Marc de Gasparo, M.D., Pharma Research Department, CIBA-GEIGY Limited, Basel CH 4002, Switzerland.

References

1. GEHLERT DR, GACKENHEIMER SL, SCHÖBER DA: Autoradiographic localization of subtypes of angiotensin II antagonist binding in the rat brain. *Neuroscience* 44:501-514, 1991
2. MENDELSON FAO, MILLAN M, QUIRION R, AGUILERA G, CHOU ST, CATT KJ: Localization of angiotensin II receptors in rat and monkey kidney by in vitro autoradiography. *Kidney Int* 31(Suppl 20):S40-S44, 1987
3. DOUGLAS JG: Angiotensin receptor subtypes of the kidney cortex. *Am J Physiol* 253:F1-F7, 1987
4. TIMMERMAN PBMWM, WONG PC, CHIU AT, HERBLIN WF, BENFIELD P, CARINI DJ, LEE RJ, WEXLER RR, SAYE JAM, SMITH RD: Angiotensin-II receptors and angiotensin-II receptor antagonists. *Pharmacol Rev* 45:205-251, 1993
5. WHITEBREAD S, MELE M, KAMBER B, DE GASPARO M: Preliminary biochemical characterization of two angiotensin II receptor subtypes. *Biochem Biophys Res Commun* 163:284-291, 1989
6. CHIU AT, HERBLIN WF, MCCALL DE, ARDECKY RJ, CARINI DJ, DUNCIA JV, PEASE LJ, WONG PC, WEXLER RR, JOHNSON AL: Identification of angiotensin II receptor subtypes. *Biochem Biophys Res Commun* 165:196-203, 1989
7. ROGGE H, SCHMID A, DE GASPARO M: Identification and characterization of angiotensin II receptor subtypes in rabbit ventricular myocardium. *Biochem Biophys Res Commun* 173:416-422, 1990
8. DE GASPARO M, WHITEBREAD S, MELE M, MOTANI AS, WHITCOMBE PJ, RAMJOUÉ HP, KAMBER B: Biochemical characterization of two angiotensin II receptor subtypes in the rat. *J Cardiovasc Pharmacol* 16(Suppl 4):S31-S35, 1990
9. GRADY EF, SECHI LA, GRIFFIN CA, SCHAMBELAN M, KALINYAK JE: Expression of AT_2 receptors in the developing rat fetus. *J Clin Invest* 88:921-933, 1991
10. EDWARDS RM, STACK EJ, WEIDLEY EF, AIYAR N, KEENAN RM, HILL DT, WEINSTOCK J: Characterization of renal angiotensin II receptors using subtype selective antagonists. *J Pharmacol Exp Ther* 260:933-938, 1992
11. BUMPUS FM, CATT KJ, CHIU AT, DE GASPARO M, GOODFRIEND T, HUSAIN A, PEACH MJ, TAYLOR DGJ, TIMMERMAN PBMWM: Nomenclature for angiotensin receptors: A report of the nomenclature committee of the council for high blood pressure research. *Hypertension* 17:720-721, 1991
12. WEIST SA, RAMPERSAUD A, ZIMMERMAN K, STEINBERG MI: Characterization of distinct angiotensin II binding sites in rat adrenal gland and bovine cerebellum using selective non-peptide antagonists. *J Cardiovasc Pharmacol* 17:177-184, 1991
13. PUCCELL AG, HODGES JC, SEN I, BUMPUS FM, HUSAIN A: Biochemical properties of the ovarian granulosa cell type 2-angiotensin II receptor. *Endocrinology* 128:1947-1959, 1991
14. BRECHLER V, LEVENS NR, DE GASPARO M, BOTTARI SP: Angiotensin AT_2 receptor mediated inhibition of particulate guanylate cyclase: A link with protein tyrosine phosphatase stimulation? *Recept Channels* 2:79-87, 1994
15. ULLIAN ME, LINAS SL: Role of receptor cycling in the regulation of angiotensin II surface receptor number and angiotensin II uptake in rat vascular smooth muscle cells. *J Clin Invest* 84:840-846, 1989
16. DUDLEY DT, HUBBELL SE, SUMMERFELT RM: Characterization of angiotensin II (AT_2) binding sites in R3T3 cells. *Mol Pharmacol* 40:360-367, 1991
17. MURPHY TJ, ALEXANDER RW, GRIENDLING KK, RUNGE MS, BERNSTEIN KE: Isolation of a cDNA encoding the vascular type-1 angiotensin II receptor. *Nature* 351:233-236, 1991
18. SASAKI K, YAMANO Y, BARDHAN S, IWAI N, MURRAY JJ, HASEGAWA M, MATSUDA Y, INAGAMI T: Cloning and expression of a complementary DNA encoding a bovine adrenal angiotensin II type-1 receptor. *Nature* 351:230-233, 1991
19. MUKOYAMA M, NAKAJIMA M, HORIUCHI M, SASAMURA H, PRATT RE, DZAU VJ: Expression cloning of type-2 angiotensin II receptor reveals a unique class of seven-transmembrane receptors. *J Biol Chem* 268:24539-24542, 1993
20. KAMBAYASHI Y, BARDHAN S, TAKAHASHI K, TSUZUKI S, INUI H, HAMAKUBO T, INAGAMI T: Molecular cloning of a novel angiotensin-II receptor isoform involved in phosphotyrosine phosphatase inhibition. *J Biol Chem* 268:24543-24546, 1993
21. BOTTARI SP, DE GASPARO M, STECKELINGS UM, LEVENS NR: Angiotensin II receptor subtypes. Characterization, signalling mechanisms and possible physiological implications. *Front Neuroendocrinol* 14:123-171, 1993
22. GAUQUELIN G, GARCIA R: Characterization of glomerular angiotensin-II receptor subtypes. *Receptor* 2:207-212, 1992
23. GIBSON RE, THORPE HH, CARTWRIGHT ME, FRANK JD, SCHORN TW, BUNTING PB, SIEGL PK: Angiotensin II receptor subtypes in renal cortex of rats and rhesus monkeys. *Am J Physiol* 261:F512-F518, 1991
24. CHANG RS, LOTTI VJ: Angiotensin receptor subtypes in rat, rabbit and monkey tissues: Relative distribution and species dependency. *Life Sci* 49:1485-1490, 1991
25. JOURDAIN M, AMIEL C, FRIEDLANDER G: Modulation of Na-H exchange activity by angiotensin II in opossum kidney cells. *Am J Physiol* 263:C1141-C1146, 1992
26. GOLDFARB DA, DIZ DI, TUBBS RR, FERRARIO CM, NOVICK AC: Angiotensin-II receptor subtypes in the human renal cortex and renal cell carcinoma. *J Urol* 151:208-213, 1994
27. DE LEON H, GARCIA R: Angiotensin II receptor subtypes in rat renal preglomerular vessels. *Receptor* 2:253-260, 1992
28. CHANSEL D, CZEKALSKI S, PHAM P, ARDAILLOU R: Characterization of angiotensin II receptor subtypes in human glomeruli and mesangial cells. *Am J Physiol* 262:F432-F441, 1992

29. CHANSEL D, VANDERMEERSCH S, PHAM P, ARDAOULLOU R: Characterization of [^3H]losartan receptors in isolated rat glomeruli. *Eur J Pharmacol-Molec Pharm* 247:193–198, 1993
30. TERADA Y, TOMITA K, NONOGUCHI H, MARUMO F: PCR Localization of angiotensin-II receptor and angiotensinogen messenger RNAs in rat kidney. *Kidney Int* 43:1251–1259, 1993
31. KAKINUMA Y, FOGO A, INAGAMI T, ICHIKAWA I: Intrarenal localization of angiotensin-II type-1 receptor messenger RNA in the rat. *Kidney Int* 43:1229–1235, 1993
32. MEISTER B, LIPPOLDT A, BUNNEMANN B, INAGAMI T, GANTEN D, FUXE K: Cellular expression of angiotensin type-1 receptor messenger RNA in the kidney. *Kidney Int* 44:331–336, 1993
33. PAXTON WG, RUNGE M, HORAIST C, COHEN C, ALEXANDER RW, BERNSTEIN KE: Immunohistochemical localization of rat angiotensin-II AT1 receptor. *Am J Physiol* 264:F989–F995, 1993
34. HERBLIN WF, DIAMOND SM, TIMMERMAN PB: Localization of angiotensin II receptor subtypes in the rabbit adrenal and kidney. *Peptides* 12:581–584, 1991
35. BERGSMAN DJ, ELLIS C, KUMAR C, NUTHULAGANTI P, KERSTEN H, ELSHOUBAGY N, GRIFFIN E, STADEL JM, AIYAR N: Cloning and characterization of a human angiotensin II type 1 receptor. *Biochem Biophys Res Commun* 183:989–995, 1992
36. FURUTA H, GUO DF, INAGAMI T: Molecular cloning and sequencing of the gene encoding human angiotensin II type 1 receptor. *Biochem Biophys Res Commun* 183:8–13, 1992
37. YE MQ, HEALY DP: Characterization of an angiotensin type-1 receptor partial cDNA from rat kidney: Evidence for a novel AT1B receptor subtype. *Biochem Biophys Res Commun* 185:204–210, 1992
38. CHIU AT, DUNSCOMB J, KOSIEROWSKI J, BURTON CRA, SANTOMENNA LD, CORJAY MH, BENFIELD P: The ligand binding signatures of the rat at(1a), at(1b) and the human at(1) receptors are essentially identical. *Biochem Biophys Res Commun* 197:440–449, 1993
39. ERNSBERGER P, ZHOU J, DAMON TH, DOUGLAS JG: Angiotensin II receptor subtypes in cultured rat renal mesangial cells. *Am J Physiol* 263:F411–F416, 1992
40. EDWARDS RM, STACK EJ: Angiotensin II inhibits glomerular adenylylate cyclase via the angiotensin II receptor subtype 1 (AT1). *J Pharmacol Exp Ther* 266:506–510, 1993
41. OLINS GM, CORPUS VM, MCMAHON EG, PALOMO MA, SCHUN JR, BLEHM DJ, HUANG HC, REITZ DB, MANNING RE, BLAINE EH: In vitro pharmacology of a nonpeptidic angiotensin II receptor antagonist, SC-51316. *J Pharmacol Exp Ther* 261:1037–1043, 1992
42. SECHI LA, GRADY EF, GRIFFIN CA, KALINYAK JE, SCHAMBELAN M: Distribution of angiotensin II receptor subtypes in rat and human kidney. *Am J Physiol* 262:F236–F240, 1992
43. TUFRO McREDDIE A, HARRISON JK, EVERETT AD, GOMEZ RA: Ontogeny of type 1 angiotensin II receptor gene expression in the rat. *J Clin Invest* 91:530–537, 1993
44. CIUFFO GM, VISWANATHAN M, SELTZER AM, TSUTSUMI K, SAAVEDRA JM: Glomerular angiotensin-II receptor subtypes during development of rat kidney. *Am J Physiol* 265:F264–F271, 1993
45. FRIBERG P, SUNDELIN B, BOHMAN SO, BOBIK A, NILSSON H, WICKMAN A, GUSTAFSSON H, PEDERSEN J, ADAMS MA: Renin-angiotensin system in neonatal rats: Induction of a renal abnormality in response to ACE inhibition or angiotensin II antagonism. *Kidney Int* 45:485–492, 1994
46. COGAN MG, LIU FY, WONG PC, TIMMERMAN PB: Comparison of inhibitory potency by nonpeptide angiotensin II receptor antagonists PD123177 and DuP 753 on proximal nephron and renal transport. *J Pharmacol Exp Ther* 259:687–691, 1991
47. WONG PC, HART SD, DUNCIA JV, TIMMERMAN PB: Nonpeptide angiotensin II receptor antagonists. Studies with DuP 753 and EXP3174 in dogs. *Eur J Pharmacol* 202:323–330, 1991
48. LOUTZENHISER R, EPSTEIN M, HAYASHI K, TAKENAKA T, FORSTER H: Characterization of the renal microvascular effects of angiotensin II antagonist DUP-753: Studies in isolated perfused hydronephrotic kidneys. *Am J Hypertens* 4:309S–314S, 1991
49. BOVEE KC, WONG PC, TIMMERMAN PB, THOOLEN MJ: Effects of the nonpeptide angiotensin II receptor antagonist DuP 753 on blood pressure and renal functions in spontaneously hypertensive PH dogs. *Am J Hypertens* 4:327S–333S, 1991
50. CLARK KL, ROBERTSON MJ, DREW GM: Effects of angiotensin AT1 or AT2 receptor blockade on basal renal function and on the renal effects of angiotensin II (Ang II) in the anaesthetized dog. *Br J Pharmacol* 106:19p, 1992
51. WOLF G, NEILSON EG: Angiotensin II induces cellular hypertrophy in cultured murine proximal tubular cells. *Am J Physiol* 259:F768–F777, 1990
52. LAFAYETTE RA, MAYER G, PARK SK, MEYER TW: Angiotensin II receptor blockade limits glomerular injury in rats with reduced renal mass. *J Clin Invest* 90:766–771, 1992
53. MENTO PF, MAITA ME, WILKES BM: Renal hemodynamics in rats with myocardial infarction: Selective antagonism of angiotensin receptor subtypes. (abstract) *Hypertension* 20:426–P66, 1992
54. MACARI D, BOTTARI S, WHITEBREAD S, DE GASPARO M, LEVENS N: Renal actions of the selective angiotensin-AT(2) receptor ligands CGP-42112B and PD-123319 in the sodium-depleted rat. *Eur J Pharmacol* 249:85–93, 1993